

# On the Maximal Energy Trees with One Maximum and One Second Maximum Degree Vertex\*

Jing Li, Xueliang Li

Center for Combinatorics and LPMC-TJKLC

Nankai University, Tianjin 300071, China

E-mail: lj02013@163.com; lxl@nankai.edu.cn

(Received June 18, 2011)

## Abstract

For a simple graph  $G$ , the energy  $E(G)$  is defined as the sum of the absolute values of all eigenvalues of its adjacent matrix. For  $d_1 > d_2 \geq 3$  and  $t \geq 3$ , denote by  $T_a$  the tree formed from a path  $P_t$  on  $t$  vertices by attaching  $d_1 - 1$   $P_2$ 's on one end and  $d_2 - 1$   $P_2$ 's on the other end of the path  $P_t$ , and  $T_b$  the tree formed from  $P_{t+2}$  by attaching  $d_1 - 1$   $P_2$ 's on an end of the  $P_{t+2}$  and  $d_2 - 2$   $P_2$ 's on the vertex next to the end. In [14] Yao showed that among trees of order  $n$  and two vertices of maximum degree  $d_1$  and second maximum degree  $d_2$  ( $d_1 > d_2$ ), the maximal energy tree is either the graph  $T_a$  or the graph  $T_b$ , where  $t = n + 4 - 2d_1 - 2d_2 \geq 3$ . However, she could not determine which one of  $T_a$  and  $T_b$  is the maximal energy tree. This is because the quasi-order method is invalid for comparing their energies. In this paper, we use a new method to determine the maximal energy tree. We prove that the maximal energy tree is  $T_b$  if  $d_1 \geq 7$ ,  $d_2 \geq 3$  or  $d_1 = 6, d_2 = 3$ . Moreover, for  $d_1 = 4$  and  $d_2 = 3$ , the maximal energy tree is the graph  $T_b$  if  $t = 4$ , and the graph  $T_a$  otherwise. For other cases, the maximal energy tree is the graph  $T_a$  if (i)  $d_1 = 5, d_2 = 4$ ,  $t$  is odd and  $3 \leq t \leq 45$ , (ii)  $d_1 = 5, d_2 = 3$ ,  $t$  is odd and  $3 \leq t \leq 29$ , (iii)  $d_1 = 6, d_2 = 5$ ,  $t = 3, 5, 7$ , (iv)  $d_1 = 6, d_2 = 4$ ,  $t = 5$ ; and for all the remaining cases, the maximal energy tree is the graph  $T_b$ .

---

\*Supported by NSFC No.11071130.

# 1 Introduction

Let  $G$  be a simple graph of order  $n$ , and  $\lambda_1, \lambda_2, \dots, \lambda_n$  be the eigenvalues of  $G$ . Then the energy of  $G$  is defined as

$$E(G) = \sum_{i=1}^n |\lambda_i|,$$

which was introduced by Gutman in [9]. The match polynomial [6, 7] of  $G$  is defined as

$$m(G, x) = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k m(G, k) x^{n-2k},$$

where  $m(G, k)$  denotes the number of  $k$ -matchings of  $G$  and  $m(G, 0) = 1$ . If  $G = T$  is a tree of order  $n$ , then the characteristic polynomial [5] of  $G$  has the form

$$\varphi(T, x) = m(T, x) = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k m(T, k) x^{n-2k}.$$

And, by Coulson integral formula [3, 4, 8, 11], we have for a tree  $T$ ,

$$E(T) = \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \log \left[ \sum_{k=0}^{\lfloor n/2 \rfloor} m(T, k) x^{2k} \right] dx.$$

As we did in [12], for convenience we use the so-called signless matching polynomial [1]

$$m^+(G, x) = \sum_{k=0}^{\lfloor n/2 \rfloor} m(G, k) x^{2k}.$$

Then we have

$$E(T) = \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \log m^+(T, x) dx. \tag{1}$$

For basic properties of  $m^+(G, x)$ , we refer to our paper [12].

For more results on graph energy, we refer to the survey [10]. For terminology and notations not defined here, we refer to the book of Bondy and Murty [2].

Graphs with extremal energies are interested in literature. In 2009 Li et al. [13] showed that among trees of order  $n$  with two vertices of maximum degree  $\Delta (\geq 3)$ , the maximal energy tree is either the graph  $G_a$  or the graph  $G_b$ , where  $t = n + 4 - 4\Delta \geq 3$  and  $G_a$  is the tree formed from a path  $P_t$  on  $t$  vertices by attaching  $\Delta - 1$   $P_2$ 's on each end of the path  $P_t$ ,  $G_b$  is the tree formed from  $P_{t+2}$  by attaching  $\Delta - 1$   $P_2$ 's on an end of the




Figure 1.1 The maximal energy trees with  $n$  vertices and two vertices  $u$  and  $v$  of degree  $d_1$  and  $d_2$ .

$P_{t+2}$  and  $\Delta - 2 P_2$ 's on the vertex next to the end. However, they could not determine which one of  $G_a$  and  $G_b$  is the maximal energy tree. In our recent paper [12], we used a new method to determine the maximal energy tree. In a similar way, Yao [14] gave the following Theorem 1.1 about the maximal energy tree with one maximum and one second maximum degree vertex.

**Theorem 1.1** ([14]) *Among trees with a fixed number of vertices ( $n$ ) and two vertices of maximum degree  $d_1$  and second maximum degree  $d_2$  ( $d_1 > d_2$ ), the maximal energy tree has as many as possible 2-branches.*

- (1) *If  $n \geq 2d_1 + 2d_2 - 1$ , then the maximal energy tree is either the graph  $T_a$  or the graph  $T_b$ , depicted in Figure 1.1.*
- (2) *If  $n \leq 2d_1 + 2d_2 - 2$ , then the maximal energy tree is among the graph  $T_c$  depicted in Figure 1.1.*

From Theorem 1.1, one can also see that for  $n \geq 2d_1 + 2d_2 - 1$ , she could not determine which one of the trees  $T_a$  and  $T_b$  has the maximal energy. In fact, the quasi-order method they used before is invalid for the special case. In this paper, we will use the Coulson integral formula method to determine which one of the trees  $T_a$  and  $T_b$  has the maximal energy. One must notice that since  $d_1 \neq d_2$  here, the energy is a function in two variables  $d_1$  and  $d_2$ , and this makes our discussion much more complicated.

## 2 Preliminaries

In this section, we list some useful properties of the signless matching polynomial  $m^+(G, x)$ , which will be used in the sequel, and already appeared in [12].

**Lemma 2.1** *Let  $v$  be a vertex of  $G$  and  $N(v) = \{v_1, v_2, \dots, v_r\}$  the set of all neighbors of  $v$  in  $G$ . Then*

$$m^+(G, x) = m^+(G - v, x) + x^2 \sum_{v_i \in N(v)} m^+(G - v - v_i, x).$$

**Lemma 2.2** *Let  $P_t$  denote a path on  $t$  vertices. Then*

$$(1) \quad m^+(P_t, x) = m^+(P_{t-1}, x) + x^2 m^+(P_{t-2}, x), \text{ for any } t \geq 1,$$

$$(2) \quad m^+(P_t, x) = (1 + x^2)m^+(P_{t-2}, x) + x^2 m^+(P_{t-3}, x), \text{ for any } t \geq 2.$$

The initials are  $m^+(P_0, x) = m^+(P_1, x) = 1$ , and we define  $m^+(P_{-1}, x) = 0$ .

**Corollary 2.3** *Let  $P_t$  be a path on  $t$  vertices. Then for any real number  $x$ ,*

$$m^+(P_{t-1}, x) \leq m^+(P_t, x) \leq (1 + x^2)m^+(P_{t-1}, x), \text{ for any } t \geq 1.$$

## 3 Main results

Before giving our main results, we state some knowledge on real analysis, for which we refer to [15].

**Lemma 3.1** *For any real number  $X > -1$ , we have*

$$\frac{X}{1+X} \leq \log(1+X) \leq X.$$

For convenience, we introduce the following notations:

$$A_1 = (x^2 + 1)(d_1 x^6 + d_2 x^6 + d_2 x^4 + d_1 d_2 x^4 + d_1 x^4 + 2x^4 + 2x^2 + d_1 x^2 + d_2 x^2 + 1),$$

$$A_2 = x^2(x^2 + 1)(x^6 + 2x^4 + d_1 d_2 x^4 + d_1 x^2 + d_2 x^2 + x^2 + 1),$$

$$B_1 = 2x^8 + d_1x^8 + 6x^6 + 2d_1d_2x^6 + d_1d_2x^4 + 2d_1x^4 + 4x^4 + 2d_2x^4 + d_2x^2 + d_1x^2 + 3x^2 + 1,$$

$$B_2 = x^2(x^2 + 1)(x^6 + 2x^4 + d_1d_2x^4 + d_1x^2 + d_2x^2 + x^2 + 1).$$

Using Lemmas 2.1 and 2.2 repeatedly, we can easily get the following two recursive formulas, where  $t = n + 4 - 2d_1 - 2d_2 \geq 3$ :

$$m^+(T_a, x) = (1 + x^2)^{d_1+d_2-5}(A_1m^+(P_{t-3}, x) + A_2m^+(P_{t-4}, x)), \tag{2}$$

and

$$m^+(T_b, x) = (1 + x^2)^{d_1+d_2-5}(B_1m^+(P_{t-3}, x) + B_2m^+(P_{t-4}, x)), \tag{3}$$

From Eqs. (2) and (3), by some elementary calculations we can obtain

$$m^+(T_a, x) - m^+(T_b, x) = (1 + x^2)^{d_1+d_2-5}(d_2 - 2)x^6(x^2 - (d_1 - 2))m^+(P_{t-3}, x). \tag{4}$$

We know directly from Figure 1.1 that if  $t = 2$  or  $d_2 = 2$ ,  $T_a \cong T_b$ , then  $E(T_a) = E(T_b)$ , so we only consider the cases  $t \geq 3$  and  $d_1 > d_2 \geq 3$ .

Now we give a useful lemma.

**Lemma 3.2** *Among trees with  $n$  vertices and two vertices of maximum and second maximum degree  $d_1$  and  $d_2$ , let  $k = d_1 - d_2$ , if  $1 \leq k \leq 3$ ,  $d_2 \geq 7 - k$  or  $4 \leq k \leq 12$ ,  $d_2 \geq 3$ , the maximal energy tree is the graph  $T_b$ , where  $t = n + 4 - 2d_1 - 2d_2 \geq 3$ .*

*Proof.* Since  $m^+(T_a, x) > 0$  and  $m^+(T_b, x) > 0$ , we have

$$\frac{m^+(T_a, x) - m^+(T_b, x)}{m^+(T_b, x)} = \frac{m^+(T_a, x)}{m^+(T_b, x)} - 1 > -1.$$

Therefore, from Eq. (1) and Lemma 3.1, we get that

$$\begin{aligned} E(T_a) - E(T_b) &= \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \log \frac{m^+(T_a, x)}{m^+(T_b, x)} dx \\ &= \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{m^+(T_a, x) - m^+(T_b, x)}{m^+(T_b, x)} \right) dx \\ &\leq \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \cdot \frac{m^+(T_a, x) - m^+(T_b, x)}{m^+(T_b, x)} dx. \end{aligned} \tag{5}$$

By Corollary 2.3, we have  $m^+(P_{t-4}, x) \leq m^+(P_{t-3}, x)$  and  $m^+(P_{t-4}, x) \geq \frac{m^+(P_{t-3}, x)}{1+x^2}$  for  $t \geq 4$ . So, we have

$$\begin{aligned} & E(T_a) - E(T_b) \\ & \leq \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \cdot \frac{m^+(T_a, x) - m^+(T_b, x)}{m^+(T_b, x)} dx \\ & = \frac{2}{\pi} \int_0^{+\infty} \frac{(d_2 - 2)x^4(x^2 - (d_1 - 2))m^+(P_{t-3}, x)}{B_1m^+(P_{t-3}, x) + B_2m^+(P_{t-4}, x)} dx \\ & \leq \frac{2}{\pi} \int_{\sqrt{d_1-2}}^{+\infty} \frac{(d_2 - 2)x^4(x^2 - (d_1 - 2))}{B_1 + B_2/(1+x^2)} dx + \frac{2}{\pi} \int_0^{\sqrt{d_1-2}} \frac{(d_2 - 2)x^4(x^2 - (d_1 - 2))}{B_1 + B_2} dx \\ & < \frac{2}{\pi} \int_{\sqrt{d_1-2}}^{+\infty} \frac{(d_2 - 2)x^4(x^2 - (d_1 - 2))}{(d_1 + 3)x^8} dx + \frac{2}{\pi} \int_1^{\sqrt{d_1-2}} \frac{(d_2 - 2)x^4(x^2 - (d_1 - 2))}{(5d_1d_2 + 6d_1 + 5d_2 + 26)x^{10}} dx \\ & + \frac{2}{\pi} \int_0^1 \frac{2(d_2 - 2)x^4(x^2 - (d_1 - 2))}{(5d_1d_2 + 6d_1 + 5d_2 + 26)(x^2 + 1)} dx = \frac{2}{\pi} f(d_1, d_2) . \end{aligned}$$

Where

$$\begin{aligned} f(d_1, d_2) & = \frac{2(d_2 - 2)}{3(d_1 + 3)\sqrt{d_1 - 2}} - \frac{d_2 - 2}{15(26 + 6d_1 + 5d_1d_2 + 5d_2)} \left( 3d_1 - 11 + \frac{2}{(d_1 - 2)^{3/2}} \right) \\ & - \frac{28d_2 - 40d_1d_2 + 80d_1 - 30\pi d_1 + 30\pi + 15\pi d_2d_1 - 56 - 15\pi d_2}{30(26 + 6d_1 + 5d_1d_2 + 5d_2)} . \end{aligned}$$

Now, for  $k = d_1 - d_2$ , we have that

- (1) if  $k = 1$ , when  $d_2 \geq 62$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (2) if  $k = 2$ , when  $d_2 \geq 60$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (3) if  $k = 3$ , when  $d_2 \geq 57$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (4) if  $k = 4$ , when  $d_2 \geq 54$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (5) if  $k = 5$ , when  $d_2 \geq 50$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (6) if  $k = 6$ , when  $d_2 \geq 47$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (7) if  $k = 7$ , when  $d_2 \geq 43$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (8) if  $k = 8$ , when  $d_2 \geq 40$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (9) if  $k = 9$ , when  $d_2 \geq 35$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .
- (10) if  $k = 10$ , when  $d_2 \geq 31$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .

(11) if  $k = 11$ , when  $d_2 \geq 24$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .

(12) if  $k = 12$ , when  $d_2 \geq 3$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} f(d_1, d_2) < 0$ .

For smaller  $d_2$ , we consider the following inequality

$$E(T_a) - E(T_b) \leq \frac{2}{\pi} \cdot g(d_1, d_2, x) < 0$$

where

$$g(d_1, d_2, x) = \int_0^{\sqrt{d_1-2}} \frac{1}{x^2} \log \left( 1 + \frac{(d_2-2)x^6(x^2 - (d_1-2))}{B_1 + B_2} \right) dx \\ + \int_{\sqrt{d_1-2}}^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{(d_2-2)x^6(x^2 - (d_1-2))}{B_1 + \frac{B_2}{1+x^2}} \right) dx .$$

By direct calculations, using a computer with the Maple programm, we can get that

(1) if  $k = 1$ , when  $6 \leq d_2 \leq 61$ ,  $E(T_a) - E(T_b) < \frac{2}{\pi} g(d_1, d_2, x) < 0$ .

(2) if  $k = 2$ , when  $5 \leq d_2 \leq 59$ ,  $E(T_a) - E(T_b) < 0$ .

(3) if  $k = 3$ , when  $4 \leq d_2 \leq 56$ ,  $E(T_a) - E(T_b) < 0$ .

(4) if  $4 \leq k \leq 11$ , when  $3 \leq d_2 \leq 53$ ,  $E(T_a) - E(T_b) < 0$ .

Then, from all the above results, we get the following conclusion: for all  $t \geq 4$ ,

(1) if  $k = 1$ , when  $d_2 \geq 6$ ,  $E(T_a) - E(T_b) < 0$ .

(2) if  $k = 2$ , when  $d_2 \geq 5$ ,  $E(T_a) - E(T_b) < 0$ .

(3) if  $k = 3$ , when  $d_2 \geq 4$ ,  $E(T_a) - E(T_b) < 0$ .

(4) if  $4 \leq k \leq 12$ , when  $d_2 \geq 3$ ,  $E(T_a) - E(T_b) < 0$ .

If  $t = 3$ , we have  $m^+(P_{t-4}, x) = m^+(P_{-1}, x) = 0$ . By a similar method as above, we can get the same result.

The proof is now complete. ■

Next we consider the case  $k \geq 13$ .

**Lemma 3.3** *Among trees with  $n$  vertices and two vertices of maximum and second maximum degree  $d_1$  and  $d_2$ , let  $k = d_1 - d_2$ , if  $k \geq 13$ ,  $d_2 \geq 3$ , then the maximal energy tree is the graph  $T_b$ , where  $t = n + 4 - 2d_1 - 2d_2 \geq 3$ .*

*Proof.* In Lemma 3.2 we proved that if  $t \geq 4, d_2 \geq 3, E(T_a) - E(T_b) < \frac{2}{\pi}f(d_1, d_2)$ . Let  $d_1 = d_2 + k$ , then  $f(d_1, d_2) = h(d_2, k)$ . We first want to show that  $h(d_2, k)$  is monotonically decreasing in  $k$ .

$$\begin{aligned} h(d_2, k) &= \frac{2(d_2 - 2)}{3(d_2 + k + 3)\sqrt{d_2 + k - 2}} \\ &\quad - \frac{d_2 - 2}{15(26 + 6(d_2 + k) + 5(d_2 + k)d_2 + 5d_2)} \left( 3(d_2 + k) - 11 + \frac{2}{(d_2 + k - 2)^{3/2}} \right) \\ &\quad - \frac{28d_2 - 40(d_2 + k)d_2 + 80(d_2 + k) - 30\pi(d_2 + k)}{30(26 + 6(d_2 + k) + 5(d_2 + k)d_2 + 5d_2)} \\ &\quad + \frac{30\pi + 15\pi d_2(d_2 + k) - 56 - 15\pi d_2}{30(26 + 6(d_2 + k) + 5(d_2 + k)d_2 + 5d_2)}. \end{aligned}$$

The derivative of  $h(d_2, k)$  on  $k$  is

$$h'(d_2, k) = h_1 + h_2 + h_3 + h_4 + h_5 + h_6,$$

where

$$\begin{aligned} h_1 &= -\frac{2(d_2 - 2)}{3(d_2 + k + 3)^2\sqrt{d_2 + k - 2}}, \\ h_2 &= -\frac{d_2 - 2}{3(d_2 + k + 3)(d_2 + k - 2)^{3/2}}, \\ h_3 &= -\frac{-30\pi - 40d_2 + 15d_2\pi + 80}{780 + 330d_2 + 180k + 150(d_2 + k)d_2}, \\ h_4 &= \frac{108d_2 - 56 - 30\pi(d_2 + k) - 40(d_2 + k)d_2 + 15d_2\pi(d_2 + k) + 30\pi - 15d_2\pi + 80k}{(780 + 330d_2 + 180k + 150(d_2 + k)d_2)^2} \\ &\quad \cdot (180 + 150d_2), \\ h_5 &= -\frac{\frac{d_2-2}{5} - \frac{d_2-2}{5(d_2+k-2)^{5/2}}}{26 + 11d_2 + 6k + 5(d_2 + k)d_2}, \\ h_6 &= \frac{\left( \frac{2}{15(d_2+k-2)^{3/2}} + \frac{3d_2+3k-11}{15} \right) (d_2 - 2)(5d_2 + 6)}{(26 + 11d_2 + 6k + 5(d_2 + k)d_2)^2}. \end{aligned}$$

Clearly,  $h_1, h_2 \leq 0$ ,

$$h_3 + h_4 = -\frac{-264d_2 - 170d_2^2 + 90d_2\pi + 75d_2^2\pi + 1208 - 480\pi}{15(5d_2^2 + 5d_2k + 11d_2 + 6k + 26)^2} < 0.$$



Moreover,

$$\begin{aligned} \frac{h_5 + h_6}{m} &= (2(d_2 + k - 2) + (3d_2 + 3k - 11)(d_2 + k - 2)^{5/2})(5d_2 + 6) \\ &\quad - 3(26 + 11d_2 + 6k + 5(d_2 + k)d_2)((d_2 + k - 2)^{5/2} - 1) \\ &= (-70d_2^3 - 140d_2^2k + 136d_2^2 - 70d_2k^2 - 8d_2k + 296d_2 - 144k^2 + 576k - 576) \\ &\quad \cdot \sqrt{d_2 + k - 2} + 25d_2^2 + 25d_2 + 25d_2k + 30k + 54 < 0, \end{aligned}$$

where

$$m = \frac{d_2 - 2}{15(d_2 + k - 2)^{5/2}(26 + 11d_2 + 6k + 5(d_2 + k)d_2)^2} > 0.$$

Thus,  $h_5 + h_6 < 0$ .

Therefore,  $h'(d_2, k) < 0$ , and hence  $h(d_2, k)$  is monotonically decreasing in  $k$ . Then, for any  $d_2 \geq 3, k \geq 13, f(d_1, d_2) = h(d_2, k) < h(d_2, 12) < 0$ . Thus  $E(T_a) - E(T_b) < 0$ .

If  $t = 3$ , we have  $m^+(P_{t-4}, x) = m^+(P_{-1}, x) = 0$ . By a similar method as above, we can get the same result. ■

From Lemmas 3.2 and 3.3, we can get the following result immediately.

**Theorem 3.4** *Among trees with  $n$  vertices and two vertices of maximum and second maximum degree  $d_1$  and  $d_2$ , if  $d_1 \geq 7$  and  $d_2 \geq 3$ , then the maximal energy tree is the graph  $T_b$ .*

Now we have proved that for most cases,  $T_b$  has the maximal energy among trees with  $n$  vertices and two vertices of maximum and second maximum degree. Only the following six special cases are left undetermined:  $(d_1, d_2) = (4, 3), (5, 4), (5, 3), (6, 5), (6, 4), (6, 3)$ . Before solving them, we give two lemmas [12] about the properties of the signless matching polynomial  $m^+(P_t, x)$  for our later use.

**Lemma 3.5** *For  $t \geq -1$ , the polynomial  $m^+(P_t, x)$  has the following form*

$$m^+(P_t, x) = \frac{1}{\sqrt{1 + 4x^2}}(\lambda_1^{t+1} - \lambda_2^{t+1}),$$

where  $\lambda_1 = \frac{1 + \sqrt{1 + 4x^2}}{2}$  and  $\lambda_2 = \frac{1 - \sqrt{1 + 4x^2}}{2}$ .

**Lemma 3.6** *Suppose  $t \geq 4$ . If  $t$  is even, then*

$$\frac{2}{1 + \sqrt{1 + 4x^2}} < \frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)} \leq 1. \quad (6)$$

*If  $t$  is odd, then*

$$\frac{1}{1 + x^2} \leq \frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)} < \frac{2}{1 + \sqrt{1 + 4x^2}}. \quad (7)$$

Note that

$$\lim_{t \rightarrow \infty} \frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)} = \frac{2}{1 + \sqrt{1 + 4x^2}}.$$

Therefore, in view of Ineq. (6), if  $t$  is even and sufficiently large, then for some  $x$ , there exists some  $\frac{2}{1 + \sqrt{1 + 4x^2}} < \Theta' < 1$ , such that  $\Theta'$  becomes an upper bound for  $\frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)}$ . Analogously, in view of Ineq. (7), if  $t$  is odd and sufficiently large, then for some  $x$  there exists some  $\frac{1}{1 + x^2} < \Theta'' < \frac{2}{1 + \sqrt{1 + 4x^2}}$ , such that  $\Theta''$  becomes a lower bound for  $\frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)}$ . By numerical testing we can find the proper  $\Theta'$  and  $\Theta''$ .

Now we are ready to deal with the case  $d_1 = 4, d_2 = 3$ .

**Theorem 3.7** *Among trees with  $n$  vertices and two vertices of maximum and second maximum degree  $d_1 = 4$  and  $d_2 = 3$ , letting  $t = n + 4 - 2d_1 - 2d_2 \geq 3$ , the maximal energy tree is the graph  $T_b$  if  $t = 4$ , and the graph  $T_a$  otherwise.*

*Proof.* By Eqs. (2), (3), (4) and (5), we have

$$\begin{aligned} E(T_a) - E(T_b) &= \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{m^+(T_a, x) - m^+(T_b, x)}{m^+(T_b, x)} \right) dx \\ &= \frac{2}{\pi} \int_0^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{(d_2 - 2)x^6(x^2 - (d_1 - 2))}{B_1 + B_2 \frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)}} \right) dx. \end{aligned} \quad (8)$$

We first consider the case that  $t$  is odd and  $t \geq 5$ . By Eq. (8) and Lemma 3.6, we have

$$\begin{aligned} &E(T_a) - E(T_b) \\ &> \frac{2}{\pi} \int_{\sqrt{2}}^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{x^6(x^2 - 2)}{B_1 + B_2 \frac{2}{1 + \sqrt{1 + 4x^2}}} \right) dx + \frac{2}{\pi} \int_0^{\sqrt{2}} \frac{1}{x^2} \log \left( 1 + \frac{x^6(x^2 - 2)}{B_1 + B_2 \frac{1}{1 + x^2}} \right) dx \\ &> \frac{2}{\pi} \cdot 0.011179 > 0. \end{aligned}$$

If  $t$  is even, we want to find  $t$  and  $x$  satisfying that

$$\frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)} < \frac{2}{-1 + \sqrt{1 + 4x^2}}. \tag{9}$$

It is equivalent to solve

$$\frac{\lambda_1^{t-3} - \lambda_2^{t-3}}{\lambda_1^{t-2} - \lambda_2^{t-2}} < -\frac{1}{\lambda_2} \quad \text{i. e.,} \quad \left( \frac{1 + \sqrt{1 + 4x^2}}{2x} \right)^{2t-6} > \sqrt{1 + 4x^2} - 1.$$

Thus,

$$2t - 6 > \log_{\frac{1 + \sqrt{1 + 4x^2}}{2x}}(\sqrt{1 + 4x^2} - 1).$$

Since for  $x \in (0, +\infty)$ ,  $\frac{1 + \sqrt{1 + 4x^2}}{2x}$  is decreasing and  $\sqrt{1 + 4x^2} - 1$  is increasing, we have that  $\log_{\frac{1 + \sqrt{1 + 4x^2}}{2x}}(\sqrt{1 + 4x^2} - 1)$  is increasing. Thus, if  $x \in [\sqrt{2}, 5]$ , then

$$\log_{\frac{1 + \sqrt{1 + 4x^2}}{2x}}(\sqrt{1 + 4x^2} - 1) \leq \log_{\frac{1 + \sqrt{101}}{10}}(\sqrt{101} - 1) < 23.$$

Therefore, when  $t \geq 15$ , i.e.,  $2t - 6 > 23$ , we have that Ineq. (9) holds for  $x \in [\sqrt{2}, 5]$ .

Now we calculate the difference of  $E(T_a)$  and  $E(T_b)$ . When  $t$  is even and  $t \geq 15$ , from Eq. (8) we have

$$\begin{aligned} & E(T_a) - E(T_b) \\ & > \frac{2}{\pi} \int_5^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{x^6(x^2 - 2)}{B_1 + B_2} \right) dx + \frac{2}{\pi} \int_{\sqrt{2}}^5 \frac{1}{x^2} \log \left( 1 + \frac{x^6(x^2 - 2)}{B_1 + B_2 \frac{2}{-1 + \sqrt{1 + 4x^2}}} \right) dx \\ & + \frac{2}{\pi} \int_0^{\sqrt{2}} \frac{1}{x^2} \log \left( 1 + \frac{x^6(x^2 - 2)}{B_1 + B_2 \frac{2}{1 + \sqrt{1 + 4x^2}}} \right) dx > \frac{2}{\pi} \cdot 0.001634 > 0. \end{aligned}$$

For  $t = 3$  and any even  $t$  with  $4 \leq t \leq 14$ , by computing the energies of the two graphs directly by a computer with Maple program, we can get that  $E(T_a) < E(T_b)$  for  $t = 4$ , and  $E(T_a) > E(T_b)$  for the other cases.

The proof is thus complete. ■

The following theorem gives the result for the cases:  $(d_1, d_2) = (5, 4), (5, 3), (6, 5), (6, 4), (6, 3)$ .

**Theorem 3.8** *Among trees with  $n$  vertices and two vertices of maximum and second maximum degree  $d_1$  and  $d_2$ , letting  $t = n + 4 - 2d_1 - 2d_2 \geq 3$ ,*

(i) for  $d_1 = 5, d_2 = 4$ , the maximal energy tree is the graph  $T_a$  if  $t$  is odd and  $3 \leq t \leq 45$ , and the graph  $T_b$  otherwise.

(ii) for  $d_1 = 5, d_2 = 3$ , the maximal energy tree is the graph  $T_a$  if  $t$  is odd and  $3 \leq t \leq 29$ , and the graph  $T_b$  otherwise.

(iii) for  $d_1 = 6, d_2 = 5$ , the maximal energy tree is the graph  $T_a$  if  $t = 3, 5, 7$ , and the graph  $T_b$  otherwise.

(iv) for  $d_1 = 6, d_2 = 4$ , the maximal energy tree is the graph  $T_a$  if  $t = 5$ , and the graph  $T_b$  otherwise.

(v) for  $d_1 = 6, d_2 = 3$ , the maximal energy tree is the graph  $T_b$  for any  $t \geq 3$ .

*Proof.* We consider the following cases separately:

(i)  $d_1 = 5, d_2 = 4$ .

If  $t$  is even, we want to find  $t$  and  $x$  satisfying that

$$\frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)} < \frac{2.1}{1 + \sqrt{1 + 4x^2}}. \tag{10}$$

It is equivalent to solve

$$2t - 6 > \log_{\frac{1 + \sqrt{1 + 4x^2}}{2x}} \left( 41 - \frac{42}{\sqrt{1 + 4x^2} + 1} \right).$$

If  $x \in [1, \sqrt{3}]$ ,

$$\log_{\frac{1 + \sqrt{1 + 4x^2}}{2x}} \left( 41 - \frac{42}{\sqrt{1 + 4x^2} + 1} \right) \leq \log_{\frac{1 + \sqrt{13}}{2\sqrt{3}}} \left( 41 - \frac{42}{1 + \sqrt{13}} \right) < 13.$$

Therefore, when  $t \geq 10$ , i.e.,  $2t - 6 > 13$ , we have that Ineq. (10) holds for  $x \in [1, \sqrt{3}]$ .

Then, if  $t$  is even and  $t \geq 10$ , from Eq. (8) and Lemma 3.6 we have

$$\begin{aligned} E(T_a) - E(T_b) &< \frac{2}{\pi} \int_{\sqrt{3}}^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{2x^6(x^2 - 3)}{B_1 + B_2 \frac{2}{1 + \sqrt{1 + 4x^2}}} \right) dx \\ &+ \frac{2}{\pi} \int_1^{\sqrt{3}} \frac{1}{x^2} \log \left( 1 + \frac{2x^6(x^2 - 3)}{B_1 + B_2 \frac{2.1}{1 + \sqrt{1 + 4x^2}}} \right) dx \\ &+ \frac{2}{\pi} \int_0^1 \frac{1}{x^2} \log \left( 1 + \frac{2x^6(x^2 - 3)}{B_1 + B_2} \right) dx < \frac{2}{\pi} \cdot (-0.000231) < 0. \end{aligned}$$

If  $t$  is odd, we want to find  $t$  and  $x$  satisfying that

$$\frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)} > \frac{1.9}{1 + \sqrt{1 + 4x^2}}, \quad (11)$$

that is

$$2t - 6 > \log_{\frac{1 + \sqrt{1 + 4x^2}}{2x}} \left( 39 - \frac{38}{\sqrt{1 + 4x^2} + 1} \right).$$

Then we get that when  $t \geq 699$ , and  $x \in [\sqrt{3}, 190]$ , the Ineq. (11) holds. Thus, if  $t$  is odd and  $t \geq 699$ , from Eq. (8) and Lemma 3.6 we have

$$\begin{aligned} & E(T_a) - E(T_b) \\ & < \frac{2}{\pi} \int_{190}^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{2x^6(x^2 - 3)}{B_1 + B_2 \frac{1}{1+x^2}} \right) dx + \frac{2}{\pi} \int_{\sqrt{3}}^{190} \frac{1}{x^2} \log \left( 1 + \frac{2x^6(x^2 - 3)}{B_1 + B_2 \frac{1.9}{1 + \sqrt{1 + 4x^2}}} \right) dx \\ & \quad + \frac{2}{\pi} \int_0^{\sqrt{3}} \frac{1}{x^2} \log \left( 1 + \frac{2x^6(x^2 - 3)}{B_1 + B_2 \frac{2}{1 + \sqrt{1 + 4x^2}}} \right) dx < \frac{2}{\pi} \cdot (-1.41 \times 10^{-5}) < 0. \end{aligned}$$

For any even  $t$  with  $4 \leq t \leq 8$  and any odd  $t$  with  $3 \leq t \leq 697$ , by computing the energies of the two graphs directly by a computer with Matlab program, we get that  $E(T_a) > E(T_b)$  for any odd  $t$  with  $3 \leq t \leq 45$ , and  $E(T_a) < E(T_b)$  for the other cases.

(ii)  $d_1 = 5, d_2 = 3$ .

If  $t$  is even and  $t \geq 4$ , from Eq. (8) and Lemma 3.6, we have

$$\begin{aligned} E(T_a) - E(T_b) & < \frac{2}{\pi} \int_{\sqrt{3}}^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{x^6(x^2 - 3)}{B_1 + B_2 \frac{2}{1 + \sqrt{1 + 4x^2}}} \right) dx \\ & \quad + \frac{2}{\pi} \int_0^{\sqrt{3}} \frac{1}{x^2} \log \left( 1 + \frac{x^6(x^2 - 3)}{B_1 + B_2} \right) dx < \frac{2}{\pi} \cdot (-1.224 \times 10^{-4}) < 0. \end{aligned}$$

If  $t$  is odd and  $t \geq 699$ , by the similar proof in (i), we get that  $E(T_a) - E(T_b) < \frac{2}{\pi} \cdot (-9.90 \times 10^{-4}) < 0$ .

For any odd  $t$  with  $3 \leq t \leq 697$ , by computing the energies of the two graphs directly with Matlab program, we get that  $E(T_a) > E(T_b)$  for any odd  $t$  with  $3 \leq t \leq 29$ , and  $E(T_a) < E(T_b)$  for the other cases.

(iii)  $d_1 = 6, d_2 = 5$ .

If  $t$  is even, by the similar method as used in (ii), we get that  $E(T_a) - E(T_b) < \frac{2}{\pi} \cdot (-0.018405) < 0$ .

If  $t$  is odd, similar to the proof in (i), we can show that when  $t \geq 27$  and  $x \in [2, 22]$ , the following inequality holds:

$$\frac{m^+(P_{t-4}, x)}{m^+(P_{t-3}, x)} > \frac{1}{1 + \sqrt{1 + 4x^2}}.$$

Hence, if  $t$  is odd and  $t \geq 27$ , we have

$$\begin{aligned} & E(T_a) - E(T_b) \\ & < \frac{2}{\pi} \int_{22}^{+\infty} \frac{1}{x^2} \log \left( 1 + \frac{3x^6(x^2 - 4)}{B_1 + B_2 \frac{1}{1+x^2}} \right) dx + \frac{2}{\pi} \int_2^{22} \frac{1}{x^2} \log \left( 1 + \frac{3x^6(x^2 - 4)}{B_1 + B_2 \frac{1}{1+\sqrt{1+4x^2}}} \right) dx \\ & \quad + \frac{2}{\pi} \int_0^2 \frac{1}{x^2} \log \left( 1 + \frac{3x^6(x^2 - 4)}{B_1 + B_2 \frac{2}{1+\sqrt{1+4x^2}}} \right) dx < \frac{2}{\pi} \cdot (-0.002914) < 0. \end{aligned}$$

For any odd  $t$  with  $3 \leq t \leq 25$ , by computing the energies of the two graphs directly, we can get that  $E(T_a) > E(T_b)$  for  $t = 3, 5, 7$ , and  $E(T_a) < E(T_b)$  for the other cases.

(iv)  $d_1 = 6, d_2 = 4$ .

If  $t$  is even, by the similar method as used in (ii), we get that  $E(T_a) - E(T_b) < \frac{2}{\pi} \cdot (-0.015171) < 0$ .

If  $t$  is odd and  $t \geq 27$ , by the similar proof in (iii), we get that  $E(T_a) - E(T_b) < \frac{2}{\pi} \cdot (-0.004557) < 0$ .

For any odd  $t$  with  $3 \leq t \leq 25$ , by computing the energies of the two graphs directly, we get that  $E(T_a) > E(T_b)$  for  $t = 5$ , and  $E(T_a) < E(T_b)$  for the other cases.

(v)  $d_1 = 6, d_2 = 3$ .

If  $t$  is even, by the similar method as used in (ii), we get that  $E(T_a) - E(T_b) < \frac{2}{\pi} \cdot (-0.009652) < 0$ .

If  $t$  is odd and  $t \geq 27$ , by the similar proof as used in (iii), we get that  $E(T_a) - E(T_b) < \frac{2}{\pi} \cdot (-0.004244) < 0$ .

For any odd  $t$  with  $3 \leq t \leq 25$ , by computing the energies of the two graphs directly, we get that  $E(T_a) < E(T_b)$  for all  $t \geq 3$ .

The proof is now complete. ■

## References

- [1] B. Lass, Matching polynomials and duality, *Combinatorica*, **24** (2004) 427–440.
- [2] J. A. Bondy, U. S. R. Murty, *Graph Theory*, Springer–Verlag, Berlin, 2008.
- [3] C. A. Coulson, On the calculation of the energy in unsaturated hydrocarbon molecules, *Proc. Cambridge Phil. Soc.* **36** (1940) 201–203.
- [4] C. A. Coulson, J. Jacobs, Conjugation across a single bond, *J. Chem. Soc.* (1949) 2805–2812.
- [5] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs – Theory and Application* Johann Ambrosius Barth Verlag, Heidelberg, 1995.
- [6] E. J. Farrell, An introduction to matching polynomials, *J. Combin. Theory Ser. B* **27** (1979) 75–86.
- [7] R. A. Beezera, E. J. Farrell, The matching polynomial of a regular graph, *Discr. Math.* **137** (1995) 7–18.
- [8] I. Gutman, Acyclic systems with extremal Hückel  $\pi$ -electron energy, *Theor. Chim. Acta (Berlin)* **45** (1977) 79–87.
- [9] I. Gutman, The energy of a graph, *Ber. Math. -Statist. Sect. Forschungszentrum Graz* **103** (1978) 1–22.
- [10] I. Gutman, X. Li, J. Zhang, Graph energy, in: M. Dehmer, F. Emmert-Streib (Eds.), *Analysis of Complex Networks: From Biology to Linguistics*, Wiley–VCH, Weinheim, 2009, pp. 145–174.
- [11] I. Gutman, O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer–Verlag, Berlin, 1986.
- [12] J. Li, X. Li, Y. Shi, On the maximal energy tree with two maximum degree vertices, *Lin. Algebra Appl.* **435** (2011) 2272–2284.
- [13] X. Li, X. Yao, J. Zhang, I. Gutman, Maximum energy trees with two maximum degree vertices, *J. Math. Chem.* **45** (2009) 962–973.
- [14] X. Yao, Maximum energy trees with one maximum and one second maximum degree vertex, *MATCH Commun. Math. Comput. Chem.* **64** (2010) 217–230.
- [15] V. A. Zorich, *Mathematical Analysis*, MCCME, Moscow, 2002.