

The Full Non-Rigid Group Theory for Trimethylamine-BH₃ Complex

M. Dabirian and A. Iranmanesh*

Department of Mathematics, Tarbiat Modarres University
P.O.Box: 14115-137, Tehran, Iran
and

Institute for Studies in Theoretical Physics and Mathematics, Tehran, Iran.
iranmana@modares.ac.ir

(Received September 6, 2004)

Abstract

A simple method is described, for calculation of character table for the symmetry groups of molecules consisting of a number XH_3 groups attached to a rigid framework. For the full non-rigid group (F-NRG) of the trimethylamine-BH₃ complex, with C_{3v} symmetry, we prove that it is a group of order 1944 with 27 conjugacy classes. The character tables of this group are calculated.

1. Introduction

The non-rigid molecule group theory (NRG) in which the dynamical symmetry operations are defined as physical operations is a new field of chemistry. Some authors in a series of papers applied this notation to determine the character table of restricted NRG of some molecules [1-7].

A molecule undergoing such large amplitude movements, between various possible configurations, is known as a non-rigid molecule. Because of this deformability, the non-rigid molecules exhibit some

* Corresponding author

interesting properties of intramolecular dynamics, which can be studied more easily resorting to Group Theory [2; p.5].

The complete set of molecular conversion operations will contain overall rotation operations, describing the molecule rotating as a whole, and intramolecular motion operations, describing molecular moieties moving with respect to the rest of the molecule. Such a set forms a group, which we call the full non-rigid group (f-NRG).

Group theory for non-rigid molecules is becoming more and more relevant, and numerous application to large amplitude vibrational spectroscopy of small organic molecules are appearing in the literature [8-16].

In [17] Longuet-Higgins investigated the symmetry groups of non-rigid molecules, where changes from one conformation to another can occur easily. In many cases, these symmetry groups are not isomorphic with any of the familiar symmetry groups of rigid molecules, and their character tables are not known. It is therefore of some interest and importance to develop simple methods of calculating these character tables, which are needed for classification of wave functions, of selection rules, etc. [18,19].

The method as described here is appropriate for molecules which consist of a number XH_3 groups attached to a rigid framework. An example of such a molecule is the trimethylamine- BH_3 complex which is considered here in some detail. The method is not appropriate in the case where the framework is linear, as in ethane. However, Bunker [20] has shown how to deal with such molecules. In computing the character table of this molecule, we use [21,22], for the standard notation and terminology on character theory.

Lomont [23] has given two method for calculating character tables. These are a satisfactory for small group, but both of them require knowledge of the class structure and hence of the group multiplication table and they become inaccessible as soon as the order of the group becomes even moderately large. For

non-rigid molecules whose symmetry group may have several thousands of elements, they are usually quite impracticable.

The alternative approach is less mechanical and is simpler in practice. It involves two steps: decomposition of the group into classes, and determination of sets of basis functions for certain representations whose characters are then determined.

2. Theory

Throughout the present paper, the concepts of direct and semi-direct products will be used. For this purpose, let us consider two groups, I and G , which have no element in common except for the identity.

If I and G are such that any element I_s commutes with any element G_r :

$$I_s G_r = G_r I_s$$

their product forms a group S which is the direct product of I and G :

$$I \times G = S$$

On the other hand, if I and G do not commute in detail but rather:

$$I G_r = G_r I$$

for all $G_r \in G$, then the group forms a set which is called the semi-direct product of I and G , and is expressed as:

$$I \wedge G = S$$

where the invariant subgroup I with respect to G is written on left side [2; p.15].

Trimethylamine-BH₃ complex occurs in its preferred conformation with three equivalent methyl groups and BH₃ group in "staggered" configuration (see figure 1).




Figure 1

At first glance, the trimethylamine-BH₃ complex presents a type of large amplitude internal motion: the three rotations of the methyl groups and rotation of BH₃. The variables which describe the three methyl rotations are $\theta_1, \theta_2, \theta_3$, and that describing the BH₃ rotation is θ_4 , the three rotation axes being the N-CH₃ bonds and fourth rotation axes being the N-BH₃ bond.

The existence of the four XH₃ groups implies the existence of 81 isoenergetic conformations, described by four equivalent C_3 non-rigid subgroups. Notice that even when the XH₃ groups are distorted because of environmental effects, the C_3 symmetry has to be respected since the hydrogen atoms are indistinguishable [24]:

$$\begin{aligned}
 C_{3_1}^I &= [\hat{E} + \hat{C}_{3_1} + \hat{C}_{3_1}^2], \\
 C_{3_2}^I &= [\hat{E} + \hat{C}_{3_2} + \hat{C}_{3_2}^2], \\
 C_{3_3}^I &= [\hat{E} + \hat{C}_{3_3} + \hat{C}_{3_3}^2],
 \end{aligned}$$

$$C'_{3_4} = [\hat{E} + \hat{C}_{3_4} + \hat{C}_{3_4}^2] \quad (1)$$

The direct product of these four subgroups contain 81 dynamical symmetry operations:

$$G_{81} = C'_{3_1} \times C'_{3_2} \times C'_{3_3} \times C'_{3_4}, \quad (2)$$

which describe 81 potential energy wells on the potential energy hypersurface.

In order to establish the remaining transformation operations, let us consider the trimethylamine-BH₃ complex in an arbitrary conformation in which $\theta_1 \neq \theta_2 \neq \theta_3 \neq \theta_4$. Since the CH₃ groups are equivalent, the rotation angles may be interchanged by a 3-fold rotation without any energy variation:

$$\begin{aligned} \hat{W} f(\theta_1, \theta_2, \theta_3, \theta_4) &\equiv f(\theta_3, \theta_1, \theta_2, \theta_4), \\ \hat{W}^2 f(\theta_1, \theta_2, \theta_3, \theta_4) &\equiv f(\theta_2, \theta_3, \theta_1, \theta_4) \end{aligned} \quad (3)$$

or by binary exchange (reflection) which induces an inversion of the rotation sense:

$$\hat{V} f(\theta_1, \theta_2, \theta_3, \theta_4) \equiv f(\theta_1, \theta_3, \theta_2, \theta_4), \quad (4)$$

$$\hat{U} f(\theta_1, \theta_2, \theta_3, \theta_4) \equiv f(\theta_2, \theta_1, \theta_3, \theta_4), \quad (5)$$

$$\hat{T} f(\theta_1, \theta_2, \theta_3, \theta_4) \equiv f(\theta_3, \theta_2, \theta_1, \theta_4), \quad (6)$$

being

$$W' = [\hat{E} + \hat{W} + \hat{W}^2],$$

$$\begin{aligned}
 V' &= [\hat{E} + \hat{V}], \\
 U' &= [\hat{E} + \hat{U}], \\
 T' &= [\hat{E} + \hat{T}]
 \end{aligned}
 \tag{7}$$

The semidirect product of these four subgroups forms another subgroup isomorphic with the C_{3v} symmetry point group:

$$G_{24} = [W' \wedge V' \wedge U' \wedge T'] \sim C_{3v},
 \tag{8}$$

which is the symmetry group of the trimethylamine-BH₃ complex, skeleton with unstructured methyl groups.

As a result, the complete f-NRG of the trimethylamine-BH₃ complex may be written as

$$G_{1944} = [W' \wedge V' \wedge U' \wedge T'] \wedge [C_{3_1}' \times C_{3_2}' \times C_{3_3}' \times C_{3_4}']
 \tag{9}$$

which is group of order 1944.

3. Determination of classes and character table

We consider the point group of the trimethylamine-BH₃ complex in the case of a rigid framework. We consider the full non-rigid group G (f-NRG) of this molecule each equilibrium conformation of which has an ordinary point-group symmetry C_{3v} .

Since G is a permutation group, every two elements of this group with different cycle structure belong to different conjugacy classes. Using figure 1, we give the cycle structure of the representatives of the conjugacy classes of G in Table 1. In this table, cc stands for conjugacy classes, n is their number, and a permutation $(\underbrace{r_1, \dots, r_a}_{b\text{-number}}) \dots (\underbrace{l_1, \dots, l_a}_{d\text{-number}}) (t_1, \dots, t_c) \dots (s_1, \dots, s_c)$ is denoted by $a^b.c^d$ for $a, b, c, d \in N$.

Table1.

cc	1	3^3	3	3^2	$2^2.3$	$2^2.3^2$	3^4	$3^2.9$	$2^3.4$	$2^3.3.4$	2^6	$2^3.6$
n	1	2	2	2	2	1	3	4	1	2	1	1

cc	3^3	3.9	2^2
n	2	2	1

We need the generator of G . For this purpose, we done the following procedure:

Let us first consider the operations which leave the framework of the molecule unchanged. In this case, each methyl group can be left alone or rotated through 120° in either direction which commute with another equivalent methyl group. We first obtain the following permutations:

a, b, c : The permutations obtained from the above rule (a obtained when the methyl group left alone and b obtained when the methyl group rotated through 120° in left direction and c obtained when the methyl group rotated through 120° in right direction) when the element of symmetry C_3 passes through the atoms: 1, 14.

d, e : The permutations obtained from the above rule (d obtained when the methyl group left alone and e obtained when the methyl group rotated through 120° in either direction) when the element of symmetry σ_v passes through the atoms: 1, 4, 11, 14,15.

f, h : The permutations obtained from the above rule (f obtained when the methyl group left alone and h obtained when the methyl group rotated through 120° in either direction) when the element of symmetry σ_v passes through the atoms: 1, 2, 5, 14, 16.

k, l : The permutations obtained from the above rule (k obtained when the methyl group left alone and l obtained when the methyl group rotated through 120° in either direction) when the element of symmetry σ_v passes through the atoms: 1, 3, 8, 14, 17.

m : The permutation obtained from the rotation of BH_3 through the symmetry C_3 passes through atoms 1 and 14.

Therefore, we can write these relations as follows:

$$a = (2,3,4)(5,8,11)(6,9,12)(7,10,13)(15,16,17);$$

$$b = (2,3,4)(5,9,13,7,8,12,6,10,11)(15,16,17);$$

$$c = (2,3,4)(5,10,12,6,8,13,7,9,11)(15,16,17);$$

$$d = (2,3)(7,9)(5,10)(6,8)(12,13)(16,17);$$

$$e = (2,3)(7,9,5,10,6,8)(12,13)(16,17);$$

$$f = (3,4)(9,13)(10,12)(8,11)(6,7)(15,17);$$

$$h = (3,4)(9,13,10,12,8,11)(6,7)(15,17);$$

$$k = (2,4)(6,13)(5,12)(7,11)(9,10)(15,16);$$

$$l = (2,4)(6,13,5,12,7,11)(9,10)(15,16);$$

$$m = (15,16,17);$$

By using the program of GAP [25], we can see that the permutation $a, b, c, d, e, f, h, k, l, m$ are generators of G (any permutation in G can be written as a product of some $a, b, c, d, e, f, h, k, l$ and m). That is,

$G = \langle a, b, c, d, e, f, h, k, l, m \rangle$. Hence G is a group of order 1944 and has 27 conjugacy classes which we are listed in Table 2.

Table 2: The Representatives of conjugacy classes of the Group G

No.	Representatives	Size
1	Identity	1
2	(5,6,7)(8,9,10)(11,12,13)	8
3	(11,12,13)	6
4	(5,7,6)(8,10,9)	12
5	(6,7)(12,13)	27
6	(5,6)(8,9,10)(11,12)	54
7	(2,3,4)(5,8,11)(6,9,12)(7,10,13)(15,16,17)	72
8	(2,3,4)(5,8,12,6,9,13,7,10,11)(15,16,17)	72
9	(2,3,4)(5,8,13,7,10,12,6,9,11)(15,16,17)	72
10	(15,17,16)	2
11	(5,6,7)(8,9,10)(11,12,13)(15,17,16)	8
12	(5,7,6)(8,10,9)(11,13,12)(15,17,16)	8
13	(11,12,13)(15,17,16)	12
14	(5,7,6)(8,10,9)(15,17,16)	24
15	(6,7)(12,13)(15,17,16)	54
16	(5,6)(8,9,10)(11,12)(15,17,16)	108
17	(2,3,4)(5,8,11)(6,9,12)(7,10,13)	72
18	(2,3,4)(5,8,12,6,9,13,7,10,11)	72
19	(2,3,4)(5,8,13,7,10,12,6,9,11)	72
20	(2,4,3)(5,11,8)(6,12,9)(7,13,10)(15,16,17)	72
21	(2,4,3)(5,12,9,6,13,10,7,11,8)(15,16,17)	72
22	(2,4,3)(5,13,10,7,12,9,6,11,8)(15,16,17)	72
23	(3,4)(8,13,10,12)(9,11)(15,17)	162
24	(3,4)(5,6,7)(8,11,10,13)(9,12)(15,17)	162
25	(3,4)(5,7,6)(8,12,10,11)(9,13)(15,17)	162
26	(3,4)(6,7)(8,12)(9,11)(10,13)(15,17)	162
27	(3,4)(6,7)(8,13,10,11,9,12)(15,17)	324

Table 3 (Continued)

	12a	12b	2b	6d
x1	1	1	1	1
x2	-1	-1	-1	-1
x3	0	0	0	0
x4	0	0	0	0
x5	0	0	0	0
x6	0	0	0	0
x7	-1	-1	1	1
x8	1	1	-1	-1
x9	0	0	0	0
x10	1	1	0	0
x11	-1	-1	0	0
x12	A	-A	0	0
x13	-A	A	0	0
x14	0	0	0	0
x15	0	0	0	0
x16	0	0	0	0
x17	0	0	0	0
x18	0	0	0	0
x19	0	0	0	0
x20	0	0	0	0
x21	0	0	0	0
x22	0	0	0	0
x23	0	0	0	0
x24	0	0	0	0
x25	0	0	-2	1
x26	0	0	2	-1
x27	0	0	0	0

4. Conclusions

The present character table for trimethylamine-BH₃ complex has been deduced from:

- the structure of group:

$$G_{1944} = [W^I \wedge V^I \wedge U^I \wedge T^I] \wedge [C_{3_1}^I \times C_{3_2}^I \times C_{3_3}^I \times C_{3_4}^I]$$

- the division of group in 27 classes, the sum of the square of the dimensions of which gives the order of the group:

$$2 \times (1)^2 + 4 \times (2)^2 + 2 \times (3)^2 + 5 \times (6)^2 + 9 \times (8)^2 + 4 \times (12)^2 + (24)^2 = 1944$$

-the composition of the 27 classes in;

1,8,6,12,27,54, 72, 72, 72, 2, 8, 8, 12, 24, 54, 108, 72, 72, 72, 72,72, 72, 162, 162, 162, 162 and 324 elements.

The potential energy function for rotation four XH_3 groups will be described by a 81-fold well potential energy hypersurface. As a result, the spectrum of the trimethylamine- BH_3 complex, is expected to possess 27 substates.

Acknowledgement. The authors wish to thank Professor I.Yavari for many useful discussions.

They are also, indebted to the referees for their valuable comment. The second author would like to thank the Institute for Studies in Theoretical Physics and Mathematics (IPM) Tehran, Iran for the financial support (No. 83200016).

References

- [1] Y. G. Smeyers and M. Villa, *J. Math. Chem.* 28(4) (2000) 377-388.
- [2] Y. G. Smeyers, *Adv. Quantum Chem.* 24 (1992) 1-77.
- [3] Y. G. Smeyers, in: *Structure and Dynamics of Non- Rigid Molecular Systems*, Y. G. Smeyers (ed), Kluwer Academic, Dordrecht. 1995, 121-151.
- [4] A. R. Ashrafi and M. Hamadani, *Croat. Chem. Acta* 76 (2003) 299-303.
- [5] M. Hamadani and A. R. Ashrafi, *Croat. Chem. Acta* 76 (2003) 305-312.
- [6] A. R. Ashrafi, *J. Appl. Math. Comput.* 14(2004) 289-303.
- [7] M. Hamadani and A. R. Ashrafi, *International J. of Mathematics and Mathematical Sciences.*, 42 (2003) 2701- 2706.
- [8] P. R. Bunker, *Mol. Phys.*, 8 (1964) 81.
- [9] J. Serre, *Intl. J. Quantum Chem.*, S 2 (1968) 107.
- [10] S. L. Altmann, “ *Induced Representation in Crystal & Molecules*”, Academic Press, London, 1977.
- [11] G. S. Ezra, “*Symmetry Properties of Molecules*”, *Lecture Note in Chemistry* 28, Springer, 1982.
- [12] J. Maruani and J. Serre (eds), “*Symmetries and Properties of Non-Rigid Molecules*”, Elsevier, Amsterdam, 1983.
- [13] Y. G. Smeyers, M. L. Senent, V. Botella and D. C. Moule, *J. Chem. Phys.*, 98 (1993) 2754.
- [14] A. Van der Avoird, *J. Chem. Phys.*, 98 (1993) 5327.
- [15] Y. G. Smeyers, M. Villa and M. L. Senent, *J. Mol. Spect.*, 191 (1998) 232.
- [16] A. Viver-Bung, V. H. Uet and Y. G. Smeyers, *J. Chem. Phys.*, 109 (1998) 2279.
- [17] H. C. Longuet-Higgins, *Mol. Phys.* 6 (1963) 445.
- [18] A. J. Stone, *J. Chem. Phys.*, 41 (1964) 1568-1579.
- [19] J. T. Hougen, *J. Mol. Spec.*, 82 (1980) 92.
- [20] P. R. Bunker, *Molecular Symmetry in Spectroscopy*, NRC Research Prss, 1998
- [21] I. M. Isaacs, *Character Theory of Finite Groups*, Academic Press, 1978.
- [22] G. James and M. Liebeck, *Representation and Characters of Goups*, Camberidge University Press, 1993.

- [23] J. S. Lomont, Applications of Finite Groups, Academic Press Inc., New York, 1959.
- [24] Y. G. Smeyers and M. Villa, Chem. Phys. Lett. 235 (1995) 576.
- [25] M. Schonert et al., GAP, Groups, Algorithms and Programming, Lehrstuhl D fuer Mathematik, RWTH, Aachen, 1992.