MATCH
Communications in Mathematical and in Computer Chemistry

MATCH Commun. Math. Comput. Chem. **97** (2027) 611–620
ISSN: 0340–6253
doi: 10.46793/match.97-2.10726

Eisenstein Integer Structure of Goldberg–Coxeter Transformations and Identification of G–C Fullerenes

Shaoqing Li[a,*]

[a] *Anhui Jianzhu University, Hefei, Anhui, China*

leesq@ahjzu.edu.cn

(Received March 25, 2026)

Abstract

Goldberg–Coxeter (G–C) transformations generate infinite families of fullerenes while preserving the global shape of their parent structures. A central problem is to determine whether a given fullerene arises from a nontrivial G–C transformation and, if so, to identify and uniquely invert the transformation. In this paper, G–C transformations are formulated within the arithmetic framework of Eisenstein integers. Each transformation corresponds to multiplication by an Eisenstein integer, and its areal scale factor coincides with the associated Eisenstein norm. Since the ring $\mathbb{Z}[\omega]$ is a unique factorization domain, every G–C transformation admits a canonical decomposition into primary sub-transformations determined by the prime factorization of its norm. We distinguish two types of Eisenstein primes: rational primes corresponding to linear k-inflations, and non-rational primes governing genuine area-expanding transformations. This arithmetic structure reduces the identification and inversion of G–C fullerenes to prime factorization and divisibility testing in $\mathbb{Z}[\omega]$, thereby providing a rigorous and practical framework for fullerene classification.

*Corresponding author.

This work is licensed under a Creative Commons "Attribution 4.0 International" license.

1 Introduction

Fullerenes are trivalent planar graphs with exactly twelve pentagonal faces. They constitute an important class of carbon structures in mathematical chemistry and combinatorial topology. Although the enumeration of fullerenes has been extensively studied, and various encoding schemes such as face-spiral representations have been developed [5, 8] (note that non-spiral fullerenes exist [1, 9]), a systematic intrinsic classification and naming scheme remains incomplete.

The Goldberg–Coxeter (G–C) transformation provides a structural construction method for fullerene graphs. Given any fullerene, one can generate an infinite family of larger fullerenes with the same three-dimensional shape via G–C transformations [4, 6]. The resulting structures are referred to as Goldberg–Coxeter fullerenes. The leapfrog transformation and k-inflation are special cases of this construction. However, intrinsic criteria for determining whether a given fullerene arises from a G–C transformation—and for recovering its parent structure—remain insufficiently understood.

A fullerene graph can be embedded in a piecewise linear manifold characterized by the surface of a master polyhedron [3, 5, 7]. The faces of this polyhedron are referred to as cells. Under a G–C transformation, the master polyhedron scales up. While the linear scale factor is generally non-integer (except in the case of k-inflation), the areal scaling factor is always an integer as it corresponds to the number of carbon atoms.

By unfolding the fullerene manifold onto the Eisenstein plane, G–C transformations can be represented as multiplication by Eisenstein integers [6]. This algebraic formulation reveals the internal structure of G–C fullerenes: Eisenstein primes correspond to elementary transformations, and the prime factorization of the scale factor determines the decomposition of composite transformations. This observation leads to a systematic method for identifying and inverting G–C fullerenes, thereby providing a foundation for fullerene classification and nomenclature.

The paper is organized as follows. Section 2 introduces the algebraic model and establishes basic identification criteria for k-inflation and

leapfrog transformations. Section 3 defines the areal scale factor and introduces primary transformations. Section 4 establishes the correspondence between prime factorization and transformation decomposition via Eisenstein primes. Section 5 presents the identification algorithm, followed by illustrative examples in Section 6 and concluding remarks in Section 7.

2 G–C transformations and basic identification

A G–C transformation replaces each triangular face in the dual graph of a fullerene (i.e., a graphene patch $(1, 0)$) with another patch (k, l), as shown in Figure 1, generating a larger structure with the same global shape.



Figure 1. The principle of the G–C transformation.

A fullerene can be unfolded onto the Eisenstein plane, which forms a hexagonal lattice in the complex plane. The six unit directions are represented by $1, \omega, \omega^2, \omega^3, \omega^4, \omega^5$, where $\omega = e^{2\pi i/6}$ is the complex root of the equation $\omega^6 = 1$. Since $\omega^3 = -1$ and $\omega^2 = \omega - 1$, every vector in the Eisenstein plane can be expressed as $a + b\omega$, with (a, b) known as Eisenstein integers. In this setting, a G–C transformation (k, l) can be expressed as multiplication by $k + l\omega$:

$$G' = (k + l\omega) \cdot G.$$

Using Eisenstein integers, this becomes

$$a' + b'\omega = (k + l\omega)(a + b\omega) = (ak - bl) + (al + bk + bl)\omega.$$

In matrix form,

$$\begin{pmatrix} a' \\ b' \end{pmatrix} = \begin{pmatrix} k & -l \\ l & k+l \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}. \tag{1}$$

If both Eisenstein integers are non-negative, they coincide with the Coxeter coordinates. This allows transformations and inversions to be computed locally on cell edges without unfolding the entire fullerene. The resulting coordinates can be converted into Coxeter form if necessary.

Equation (1) also provides an identification criterion: if a pair (k, l) exists such that the inverse transformation yields integer coordinates, then the fullerene is a G–C fullerene, and the result corresponds to its parent structure. To carry out such identification, we should first obtain reasonable candidates for the G–C transformation.

It is not necessary to calculate all the cell edges of a fullerene for transformation, inversion, or identification. A fullerene has 20 dimensions [7]. The Coxeter coordinates for 10 independent cell edges fully determine the structure.

Theorem 1. *A fullerene is a k-inflation fullerene if and only if all Coxeter coordinates are divisible by an integer k.*

Proof. According to Equation (1), if $l = 0$, then

$$\begin{pmatrix} a' \\ b' \end{pmatrix} = \begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} ka \\ kb \end{pmatrix}. \tag{2}$$

Thus, the lattice is scaled uniformly. Such a G–C transformation is called k-inflation. Conversely, if the Coxeter coordinates of all cell edges share a common factor $k > 1$, dividing by k yields integer coordinates for the parent structure. ■

Theorem 2. *A fullerene arises from a leapfrog transformation if and only if each pair of Coxeter coordinates of the cell edges are congruent modulo* 3.

Proof. The G–C transformation $(1, 1)$ is also called the leapfrog transfor-

mation. According to Equation (1), we can obtain

$$\begin{pmatrix} a' \\ b' \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix}.$$

Solving for (a, b) in terms of (a', b') gives

$$\begin{pmatrix} a \\ b \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 2 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} a' \\ b' \end{pmatrix}. \tag{3}$$

Thus (a, b) are integers if and only if $a' \equiv b' \pmod 3$. ■

3 Areal scale factor and primary transformations

The number of carbon atoms is used as the equivalent area of a graphene patch, with the patch $(1, 0)$ assigned area 1 because the patch $(1, 0)$ contains exactly one carbon atom in the dual representation. For a triangular patch spanned by vectors $a + b\omega$ and $c + d\omega$, the area is given by the absolute value of the determinant

$$A = \begin{vmatrix} a & b \\ c & d \end{vmatrix}.$$

In particular, the area of patch (k, l) in Figure 1, spanned by the vectors $k + l\omega$ and $l + (k + l)\omega$, equals the norm of the edge:

$$N = \begin{vmatrix} k & -l \\ l & k + l \end{vmatrix} = k^2 + kl + l^2. \tag{4}$$

Thus, N is the areal scale factor of the transformation.

Theorem 3. *For a G–C fullerene, the norms of its cell edges are all integers sharing a common factor equal to the areal scale factor of the G–C transformation.*

Proof. According to equation (4), the norms of cell edges are all integers.

With the G–C transformation, they scale up with the areal scale factor. Therefore, the areal scale factor is the common factor of the norms of cell edges. ■

Using the areal scale factor, we can derive candidates for the G–C transformation via (4). Table 1 lists transformations with scale factors $N < 50$. For brevity, the transformation (k, l) in the table also represents the chiral one (if present), which has the same areal scale factor.

As shown in Table 1, many G–C transformations can be decomposed into sequences of simple transformations. There are two kinds of basic transformations: k-inflations and primary G–C transformations. A G–C transformation is called *primary* if it is neither a k-inflation nor decomposable into simpler transformations. The areal scale factors of the primary G–C transformations in Table 1 are all prime numbers.

Table 1. The G–C transformations (k, l) and their area scale factors $N < 50$; Chiral counterparts (l, k) are omitted for brevity.

(k,l)	N	Notes	(k,l)	N	Notes
$(1,1)$	3	*Primary*	$(3,3)$	27	$(1,1)\cdot(3,0)$
$(2,0)$	4	*Inflation*	$(4,2)$	28	$(2,0)\cdot(2,1)$
$(2,1)$	7	*Primary*	$(5,1)$	31	*Primary*
$(3,0)$	9	*Inflation*	$(6,0)$	36	*Inflation*
$(2,2)$	12	$(1,1)\cdot(2,0)$	$(4,3)$	37	*Primary*
$(3,1)$	13	*Primary*	$(5,2)$	39	$(1,1)\cdot(3,1)$
$(4,0)$	16	*Inflation*	$(6,1)$	43	*Primary*
$(3,2)$	19	*Primary*	$(4,4)$	48	$(1,1)\cdot(4,0)$
$(4,1)$	21	$(1,1)\cdot(2,1)$	$(5,3)$	49	$(2,1)\cdot(2,1)$
$(5,0)$	25	*Inflation*	$(7,0)$	49	*Inflation*

4 Eisenstein primes

The arithmetic of G–C transformations is governed by the structure of the Eisenstein integer ring $\mathbb{Z}[\omega]$, which is a unique factorization domain. Consequently, every nonzero Eisenstein integer admits a factorization into irreducible elements, known as Eisenstein primes, unique up to multiplication by units and ordering. There are two types of Eisenstein primes. For

the G–C transformations $k+l\omega$, as k and l are all nonnegative integers, an Eisenstein integer z is prime (up to multiplication by a unit) if and only if one of the following holds:

1. z is a natural prime p with $p \equiv 2 \pmod 3$;
2. $|z|^2 = p$, where p is a natural prime $p \equiv 0$ or $1 \pmod 3$.

In the context of G–C transformations, these two types of Eisenstein primes correspond naturally to two types of elementary transformations:

- **Type 1** (rational primes) correspond to k-inflations, arising from primes $p \equiv 2 \pmod 3$; in this case $N(z) = p^2$;
- **Type 2** (non-rational primes) correspond to primary G–C transformations, whose areal scale factors are given by primes $p = 3$ or $p \equiv 1 \pmod 3$; in this case $N(z) = p$.

Thus, the decomposition of a G–C transformation into elementary transformations is directly governed by the prime factorization of its areal scale factor in $\mathbb{Z}[\omega]$. So, the areal scale factors of the primary G–C transformations are all prime numbers, as mentioned above. Therefore, we have the following theorem.

Theorem 4. *For a fullerene without k-inflation, each prime factor of the area scale factor corresponds to a primary sub-transformation.*

Remark. The decomposition of a G–C transformation is unique up to ordering of commuting factors, mirroring the uniqueness of prime factorization in $\mathbb{Z}[\omega]$.

5 Identification algorithm

Based on the above analysis, a G–C fullerene can be identified via the following procedure.

Step 1 – Remove inflation. Compute the greatest common divisor k of all Coxeter coordinates of the cell edges. If $k > 1$, the fullerene is

a k-inflation. Divide all coordinates by k to obtain a reduced fullerene without k-inflation.

Step 2 – Compute area scale factor. For the reduced fullerene, compute the area scale factor N as the greatest common divisor of the norms $a^2 + ab + b^2$ of all cell edges. If $N = 1$, the reduced fullerene is a base fullerene (not a G–C transform of any smaller fullerene). If $N > 1$, proceed to Step 3.

Step 3 – Prime factorization. Factor N into rational primes. Each prime factor p corresponds to a primary transformation. Retrieve the parameters (k, l) from a correspondence table (e.g., Table 1, extended as needed). Two chiral candidates (k, l) and (l, k) may exist.

Step 4 – Verification and recursion. For each chiral candidate, construct the corresponding transformation sequence and invert the reduced coordinates according to Equation (1). If integer coordinates result, the candidate is valid.

In this way, the G–C transformation, including its sub-transformations, is identified, and the ultimate parent structure of the fullerene is obtained.

The procedure is computationally efficient, as it relies only on integer arithmetic and GCD computations.

6 Examples

Example 1 (Leapfrog)**.** Consider fullerene C_{60}. Its cell-edge coordinates satisfy $a \equiv b \pmod 3$. The GCD of all coordinates is $k = 1$ (no inflation). The common norm is $N = 3$, corresponding to primary transformation $(1, 1)$ (leapfrog). Inversion yields C_{20}, whose coordinates have $N = 1$. Thus, C_{60} is identified as a leapfrog transform of C_{20}.

Example 2 (Composite)**.** Consider a fullerene generated by $(4, 2)$. Its coordinates have $N = 28 = 2^2 \times 7$. The squared factor 2 corresponds to a 2-inflation, consistent with the $N(z) = p^2$ relation for Type 1 Eisenstein primes discussed in Section 4; the factor 7 corresponds to the primary transformation $(2, 1)$. The decomposition is $(4, 2) = (2, 0) \cdot (2, 1)$. Inversion recovers the parent structure.

7 Conclusion

This paper develops an arithmetic framework for Goldberg–Coxeter transformations based on the ring of Eisenstein integers. By identifying each G–C transformation with multiplication by $k + l\omega$, we show that the areal scale factor coincides with the Eisenstein norm $k^2 + kl + l^2$. The unique factorization property of $\mathbb{Z}[\omega]$ yields a canonical decomposition of any G–C transformation into elementary k-inflations and primary transformations, corresponding directly to the prime factorization of the norm. This establishes a natural correspondence between algebraic number theory and the geometric structure of fullerenes.

Building on this structure, we derive an efficient procedure for determining whether a given fullerene arises from a G–C transformation and for recovering its parent structure. The method relies solely on integer arithmetic on Coxeter coordinates and requires no global unfolding, allowing all computations to be performed locally on cell edges.

By identifying base fullerenes (those with $N = 1$) and the sequence of transformations leading to any G–C fullerene, the results provide a systematic foundation for the classification and nomenclature of fullerenes. To our knowledge, this constitutes the first purely arithmetic characterization of G–C transformations.

References

[1] G. Brinkmann, J. Goedgebeur, B. D. McKay, The smallest fullerene without a spiral, *Chem. Phys. Lett.* **522** (2012) 54–55.

[2] H. S. M. Coxeter, Virus macromolecules and geodesic domes, in: J. C. Butcher (Ed.), *A Spectrum of Mathematics: Essays Presented to H. G. Forder*, Oxford Univ. Press, Oxford, 1971, pp. 98–107.

[3] M. Deza, V. Grishukhin, Face-regular bifaced polyhedra, *J. Stat. Plan. Inference* **95** (2001) 175–195.

[4] M. Dutour Sikirić, M. Deza, Goldberg–Coxeter construction for 3- or 4-valent plane graphs, *El. J. Comb.* **11** (2004) #R20.

[5] P. W. Fowler, D. E. Manolopoulos, *An Atlas of Fullerenes*, Dover Pub., Mineola, NY, 2006.

[6] M. Goldberg, A class of multi-symmetric polyhedra, *Tohoku Math. J.* **43** (1937) 104–108.

[7] S. Li, On the intrinsic symmetries and parameterization of fullerenes, *Comput. Theor. Chem.* **1200** (2021) #113212.

[8] D. E. Manolopoulos, P. W. Fowler, A fullerene without a spiral, *Chem. Phys. Lett.* **204** (1993) 1–7.

[9] L. N. Wirz, J. E. Avery, P. Schwerdtfeger, Structure and properties of the non-face-spiral fullerenes T-C_{380}, D_3-C_{384}, D_3-C_{440} and D_3-C_{672} and their halma and leapfrog transforms, *J. Chem. Inf. Model.* **54** (2014) 121–130.